

CALIFORNIA INSTITUTE OF TECHNOLOGY
Control and Dynamical Systems

CDS 202

Problem Set #5

R. Murray
Winter 2009

Issued: 6 Feb 09 (Fri)
Due: 13 Feb 09 (Fri)

Reading: Abraham, Marsden, and Ratiu (MTA), section 4.2, 4.4

Problems:

1. Consider the following vector fields on \mathbb{R}^3 :

$$X(x) = \frac{\partial}{\partial x_2} - x_1 \frac{\partial}{\partial x_3} \quad Y(x) = \frac{\partial}{\partial x_1}.$$


Let $x_0 = (0, 0, 0)$. Show that $\phi_h^{-Y} \circ \phi_h^{-X} \circ \phi_h^Y \circ \phi_h^X(x_0) = h^2 \phi^{[X,Y]}(x_0)$.

2. Show that if Δ is a distribution of the form

$$\Delta = \text{span}\{X_1, \dots, X_d\}$$

and we have $[X_i, X_j] \in \Delta$ for all i, j then for *any* $X, Y \in \Delta$, $[X, Y] \in \Delta$. That is, to check involutivity of a distribution, we need only check that the pairwise brackets between basis elements lie in the distribution.

3. [Boothby, page 164, #4] Let $N \subset M$ be a submanifold and let $X, Y \in \mathcal{X}(M)$ be vector fields such that $X_p, Y_p \in T_p N$ for $p \in N$. Show that $[X, Y]_p \in T_p N$ for all $p \in N$.
4. [Boothby, page 164, #5] Let $F : M \rightarrow N$ be a smooth submersion of M onto N . Show that $F^{-1}(q)$ for all $q \in N$ are the leaves of a foliation on M .
5. Consider the motion of a disk rolling on the plane, as shown below:



We can represent the configuration of the disk by the xy location of the disk and the angle of the disk with respect to a fixed line on the plane. We ignore the angle through which the disk rolls. We model the motion of the disk using a one vector field to represent the *drive* input (rolling) and another vector field to represent the *steer* input (twisting).

Let $M = \mathbb{R}^2 \times S^1$ and let $p = (x, y, \theta)$ represent a point on M . Consider the two vector fields

$$X(p) = \cos \theta \frac{\partial}{\partial x} + \sin \theta \frac{\partial}{\partial y} \quad Y(p) = \frac{\partial}{\partial \theta}$$

(X is the drive vector field and Y is the steer vector field.)

- (a) Compute the flows of X and Y . Are they complete?
- (b) Verify that X and Y are invariant under their own flows.
- (c) Compute the Lie bracket between X and Y . Show that the tangent vectors X_p , Y_p , and $[X, Y]_p$ span T_pM for all $p \in M$.
- (d) Consider the change of coordinates $z = \phi(x)$ given by

$$[z_1 \quad z_2 \quad z_3] = \begin{bmatrix} x \cos \theta + y \sin \theta \\ x \sin \theta - y \cos \theta \\ \theta \end{bmatrix}.$$

(The new coordinates have the physical interpretation of being the origin of the spatial reference frame when viewed from a coordinate frame attached to the disk).

Using the pushforward map for $\phi : \mathbb{R}^3 \rightarrow \mathbb{R}^3$, compute the X and Y vector fields in this new set of coordinates. (Hint: the final answer has a pretty simple form.)

- (e) Show that $\phi_*[X, Y] = [\phi_*X, \phi_*Y]$ by explicit calculation.
- (f) Show that X and Y are invariant under the group of rigid motions on \mathbb{R}^2 , given by mappings

$$F(x, y, \theta) = \begin{bmatrix} x \cos \alpha - y \sin \alpha + \beta \\ x \sin \alpha + y \cos \alpha + \gamma \\ \theta + \alpha \end{bmatrix},$$

where $\alpha, \beta, \gamma \in \mathbb{R}$ are arbitrary constants.