

CALIFORNIA INSTITUTE OF TECHNOLOGY
Bioengineering and Biology

Bi/BE 250c

M. Elowitz and R. M. Murray
Winter 2011

Problem Set #1

Issued: Jan 6
Due: Jan 13

1. Consider a cascade of three activators $X \rightarrow Y \rightarrow Z$. Protein X is initially present in the cell in its inactive form. The input signal of X, S_x , appears at time $t=0$. As a result, X rapidly becomes active and binds the promoter of gene Y, so that protein Y starts to be produced at rate β . When Y levels exceed a threshold K , gene Z begins to be transcribed and translated at rate γ . All proteins have the same degradation/dilution rate α .
 - a) What are the concentrations of proteins Y and Z as a function of time?
 - b) What is the minimum duration of the pulse S_x such that Z will be produced?
 - c) What is response time of protein Z with respect to the time of addition of S_x ?
2. Consider a positive transcriptional feedback loop composed of two negative interactions $X \dashv Y$ and $Y \dashv X$.
 - a) Write the ODEs for the system above using Hill functions for the repressors. Assume that the two transcription/repression mechanisms have the same dynamics and both genes are degraded at the same rate 0.2. Let the basal transcription rate be 1, $K=2$, $n=2$.
 - b) To solve for the steady states, plot the *nullclines* by solving $\frac{dX}{dt} = 0$ and $\frac{dY}{dt} = 0$ (i.e. solve for $Y = g_1(X)$ where $\frac{dX}{dt} = 0$ and $Y = g_2(X)$ where $\frac{dY}{dt} = 0$ and plot both solutions). The steady states are given by the intersections of the two nullclines.
 - c) Plot the time response of X and Y using the following two initial conditions: $(X(0), Y(0)) = (1, 4)$ and $(4, 1)$. Next, plot the phase plane of the system using *pplane* in MATLAB. How do the responses change with initial conditions? Describe a situation where this type of interaction would be useful.
3. Consider the following network $X \rightarrow Y$ and $X \rightarrow X$.
 - a) Write the ODEs for the system above. Use basal expression $\beta_X = \beta_Y = 2$ and activation coefficients $K_X = 1$, $K_Y = 2$, $n_1 = n_2 = 2$. The degradation coefficients for X and Y are both 0.5.
 - b) Plot the vector field using *pplane*. How many steady states do you observe?
 - c) Solve for the steady states of the system using the derived ODEs, linearize the system and do a stability analysis.